

Specimen Label

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions.

IMPORTANT: Do not expose children, pets, or other nontarget animals to rodenticides. To help to prevent accidents:

1. Store product not in use in a location out of reach of children and pets.
2. Apply bait in locations out of reach of children, pets, domestic animals and nontarget wildlife, or in tamper-resistant bait stations. These stations must be resistant to destruction by dogs and by children under six years of age, and must be used in a manner that prevents such children from reaching into bait compartments and obtaining bait. If bait can be shaken from stations when they are lifted, units must be secured or otherwise immobilized. Even stronger bait stations are needed in areas open to hoofed livestock, raccoons, bears, or other potentially destructive animals, or in areas prone to vandalism.
3. Dispose of product container and unused, spoiled, and unconsumed bait as specified on this label.

USE RESTRICTIONS: For control of house mice in and around homes, industrial, commercial and agricultural buildings, and similar man-made structures. This product may also be used inside transport vehicles (ships, trains, aircraft) and in related port or terminal buildings.

SELECTION OF TREATMENT AREAS: Determine areas where mice will most likely find and consume the bait. Generally, these are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been seen. Protect bait from rain or snow. Remove as much alternative food as possible.

APPLICATION DIRECTIONS

Apply 1/4 to 1/2 oz. (1-2 level tablespoons) of bait at each placement location. Space placements at 8- to 12- foot intervals. Larger placements (up to 1 ounce) may be needed at

Quintox®

MOUSE SEED

KILLS MICE

ACTIVE INGREDIENT: Cholecalciferol 0.075%
INERT INGREDIENTS 99.925%
TOTAL 100.000%

KEEP OUT OF REACH OF CHILDREN CAUTION

(See right for additional precautionary statements)

Mfg. by:



Bell Laboratories, Inc.
Madison, WI 53704

EPA EST. NO. 12455-WI-1 EPA REG. NO. 12455-57

points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 10 days or until signs of mouse activity cease.

Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and leftover bait properly. To prevent reinfestation, limit sources of rodent food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present establish permanent bait stations and replenish as needed.

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION**

Keep away from humans, domestic animals, and pets.

STATEMENT OF PRACTICAL TREATMENT
If swallowed, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Avoid use of all oils.

NOTICE TO PHYSICIAN: If serum calcium levels are elevated, treatment with Calcitonin is effective in reducing calcium to normal levels. Continue monitoring serum calcium and treat as necessary for hypercalcemia (Reference, AMA Drug Evaluations, Third Edition (1977), Chapter 16, pp. 248-251).

ENVIRONMENTAL HAZARDS
Keep out of lakes, streams or ponds.

STORAGE AND DISPOSAL
Do not contaminate water, food, or feed by storage or disposal.
STORAGE: Store only in original container, in a dry area inaccessible to children and pets.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty pail. Then dispose of empty pail in sanitary landfill or by incineration or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.